

Rehabilitation Following Revision Unilateral Patellar Tendon Repair

I. <u>Immediate Postoperative Phase (Days 1-7)</u>

Goals: Restore full passive knee extension

Diminish pain and joint swelling Restore patellar mobility Initiate early controlled motion *Controlled forces on repair site

Postoperative Days 1-4

Brace: Drop locked brace locked at 0° extension with compression wrap

Sleep in brace, ambulate in brace

Weight bearing: Toe touch-25% BW WB with crutches, brace in full extension

Range of motion: 0-20°

Exercises: Ankle pumps

Quad sets

Patellar mobilization

• Ice and elevation: 20 minutes of each hour at elevate

Postoperative Day 4-7

Brace: Continue use of brace (sleep in brace locked)

Weight bearing: Locked brace two crutches, continue toe-touch-25% WB

• Range of motion: 0-30°

No CPM

Exercises: Continue all listed exercises

• Ice and elevation: Continue ice and elevation

II. <u>Maximum Protection Phase (Weeks 2 to 6)</u>

Goals: Control forces on healing tendon

Gradually increase knee flexion Restore full passive knee extension

Restore patellar mobility Retard muscular atrophy

Week 2:

• Brace: Continue use of locked brace (4-6 weeks)

Sleep in brace (4-6 weeks)

Weight bearing: Toe Touch WB (usually <25% BW)



PROM knee flexion only 0-30°
Full passive knee extension
Patellar mobilization

• Exercises: Electrical muscle stimulation to quads

Quad sets Ankle pumps

Ice and elevation: Continue ice, elevation and compression

Week 3:

• Exercise: Continue above mentioned exercises

• Range of Motion: ROM: 0-45°

Continue use of two crutches – 50% BW with locked brace

Week 4:

• Exercise: Continue all exercises listed above

• Weight bearing: 50% with 2 crutches

Range of Motion: PROM: 0-60°

Exercises: Continue quads sets & gravity eliminated SLR

Weeks 5-6:

Discontinue use of 1 crutch at 5 weeks postop

• Brace: Unlock brace for ambulation at 6 weeks

Range of motion: PROM

Week 5: 0-75° Week 6: 0-75°

• Exercises: Initiate pool exercises

Continue all exercises listed above

• Ice and elevation: Continue ice, compression and elevation as needed

III. Moderate Protection Phase (Weeks 7-16)

Goals: Control forces during ambulation and ADL's

Progress knee flexion ROM

Improve lower extremity muscular strength Restore limb confidence and function

Weeks 7-10:

Brace: Use postop brace unlocked for ambulation until 7-8 weeks

(or until determined safe)
Discontinue sleeping in brace



Range of motion: Week 7: 0-80°

Week 8: 0-90° Week 10: 0-100°

Exercises: *Gradually increase muscular strength

Straight leg raises (flexion)

Hip abd/adduction

Bicycle Pool program

Weeks 12-16:

• Range of motion: Week 12: 0-110°

• Exercises: *Continue all exercises listed above

Initiate light leg press Initiate lateral step ups

*No sports!

IV. <u>Light Activity Phase (Months 4-6)</u>

Goals: Enhancement of strength, endurance

Initiate functional activities

Improve tensile strength properties of tendon

• Exercises: Continue strengthening exercises

Continue emphasis on eccentrics and functional exercises May begin to gradually increase knee flexion beyond 115

V. Return to Sport Activity Phase (Months 7-9)

Criteria to Return to Activities:

- 1. Satisfactory clinical exam
- 2. Satisfactory strength
- 3. Proprioception test 80% > of opposite leg
- 4. Appropriate rehab progression

Goals: Gradual return to full-unrestricted sports

Continue strength program Continue functional/agility drills

Exercises: Continue strengthening exercises

Continue agility program*

Continue sport specific training*

^{*} Physician decision regarding activities and when to initiate